

BstDE I **(Dde I)**


Product No : RE1212
Quantity : 500u


Lot :
 Expiry Date :
 Concentration : 20u/μl
 Supplied with : 1ml of 10X Buffer V2
 1ml of 10X Buffer UB
 0.5ml Diluent Viva Buffer A
 (BSA included in all Reaction Buffer)

Store at -20°C


info@vivantechnologies.com
Reaction Conditions:
Buffer V2,

 10mM Tris-HCl (pH 7.5 at 30°C), 10mM MgCl₂,
 50mM NaCl, and 100μg/ml BSA.

Incubate at 60°C.
Dilution: Viva Buffer A

 10mM Tris-HCl (pH 7.4 at 25°C), 50mM KCl, 0.1mM EDTA,
 1mM DTT, 200μg/ml BSA and 50% glycerol.

Thermal Inactivation: None

Storage Buffer:


 10mM KH₂PO₄ (pH 7.5), 50mM KCl, 0.1mM EDTA,
 7mM 2-mercaptoethanol, 200μg/ml BSA and 50% glycerol.

Unit Definition:

 1u is defined as the amount of enzyme that is required to digest 1μg
 of DNA in 1 hour at 60°C in 50μl of assay buffer.

Quality Control Assays:
Ligation/ Recutting Assay:

 After 20-fold overdigestion with **BstDE I**, 90% of
 the DNA fragments can be ligated and recut.

 λ DNA
 1.2% Agarose

Overdigestion assay:

 An unaltered banding pattern was observed after 1μg
 of DNA was digested with 40u of **BstDE I** for 16
 hours at 60°C.

Activity in Reaction Buffer				
V1	V2	V3	V4	V5
75%	100%	50%	75%	50%

Buffer UB			
0.5X	1.0X	1.5X	2.0X
75%	50%	75%	50%

* Buffer UB is provided for double digestion purpose.

NOTE:

- * Total reaction volume dependent on experiment.
- * The amount of enzyme to be used is very much dependent on the DNA template.
- * For plasmid DNA, 5-10X more enzyme is required.

Example of Digestion Reaction

Enzyme : 1 unit
 Lambda 0.3μg/μl : 3.33μl (1μg DNA)
 10X Reaction Buffer : 5μl
 Sterile Distilled Water : Up to 50μl

Product Use Limitation

 This product is for research purposes and *in vitro* use only.