

### *Eco*ICR I (Sac I\*)



Product No : RE1258

Quantity : 200u



Lot :  
 Expiry Date :  
 Concentration : 10u/μl  
 Supplied with : 1ml of 10X Buffer V2  
 1ml of 10X Buffer UB  
 0.5ml Diluent Viva Buffer A

(BSA included in all Reaction Buffer)

Store at -20°C



info@vivantechnologies.com

#### Reaction Conditions:

##### Buffer V2 ,

10mM Tris-HCl (pH 7.5 at 30°C), 10mM MgCl<sub>2</sub>,  
 50mM NaCl, and 100μg/ml BSA .

**Incubate at 37°C.**

##### Dilution: Viva Buffer A

10mM Tris-HCl (pH 7.4 at 25°C), 50mM KCl, 0.1mM EDTA,  
 1mM DTT, 200μg/ml BSA and 50% glycerol.

**Thermal Inactivation:** 65°C for 20 minutes

##### Storage Buffer:

10mM Tris-HCl (pH 7.5), 200mM NaCl, 0.1mM EDTA,  
 7mM 2-mercaptoethanol, 200μg/ml BSA and 50% glycerol.

##### Unit Definition:


1u is defined as the amount of enzyme that is required to digest  
 1μg of DNA in 1 hour at 37°C in 50μl of assay buffer.

#### Quality Control Assays:

##### Ligation/ Recutting Assay:

After 10-fold overdigestion with *Eco*ICR I, more than  
 95% of the DNA fragments can be ligated and recut.

λ DNA  
 (Hind III Digest)  
 1.0% Agarose



##### Overdigestion assay:

An unaltered banding pattern was observed after 1μg of  
 DNA was digested with 20u of *Eco*ICR I for 16 hours at  
 37°C.

Activity in Reaction Buffer				
V1	V2	V3	V4	V5
100%	100%	50%	75%	75%

Buffer UB			
0.5X	1.0X	1.5X	2.0X
100%	100%	75%	75%

\* Buffer UB is provided for double digestion purpose.

#### NOTE:

- \* Total reaction volume dependent on experiment.
- \* The amount of enzyme to be used is very much dependent on the DNA template.
- \* For plasmid DNA, 5-10X more enzyme is required.

#### Example of Digestion Reaction

Enzyme : 1 unit  
 Lambda (*Hind* III Digest) 0.3μg/μl : 3.33μl (1μg DNA)  
 10X Reaction Buffer : 5μl  
 Sterile Distilled Water : Up to 50μl

#### Product Use Limitation

This product is for research purposes and *in vitro* use only.