



## Ama87 I (Ava I)



Product No. : RE1122  
Quantity : 600u



Lot :  
Expiry Date :  
Concentration : 10u/μl  
Supplied with : 1ml of 10X Buffer Ama87 I  
1ml of 10X Buffer UB  
0.5ml Diluent Viva Buffer A  
(BSA included in all Reaction Buffer)

Store at -20°C



info@vivantechnologies.com

### Reaction Conditions:

#### Buffer Ama87 I,

20mM Tris-HCl (pH 8.5), 10mM MgCl<sub>2</sub>,  
200mM NaCl and 1mM DTT.

**Incubate at 37°C.**

#### Dilution: Viva Buffer A

10mM Tris-HCl (pH 7.4 at 25°C), 50mM KCl, 0.1mM EDTA,  
1mM DTT, 200μg/ml BSA and 50% glycerol.

**Thermal Inactivation:** 65°C for 20 minutes

#### Storage Buffer:

10mM KH<sub>2</sub>PO<sub>4</sub> (pH 7.2), 100mM NaCl, 0.1mM EDTA,  
7mM 2-mercaptoethanol and 50% glycerol.

#### Unit Definition:

1u is defined as the amount of enzyme that is required to digest  
1μg of DNA in 1 hour at 37°C in 50μl of assay buffer.

#### Quality Control Assays:


##### Ligation/ Recutting Assay:

After 10-fold overdigestion with **Ama87 I**, more  
than 90% of the DNA fragments can be ligated  
and recut.

##### Overdigestion assay:

An unaltered banding pattern was observed after 1μg  
of DNA was digested with 20u of **Ama87 I** for 16  
hours at 37°C.

λ DNA  
0.7% Agarose



Activity in Reaction Buffer				
V1	V2	V3	V4	V5
10%	25%	50%	75%	10%

Buffer UB			
0.5X	1.0X	1.5X	2.0X
25%	50%	50%	75%

\* Buffer UB is provided for double digestion purpose.

#### NOTE:

- \* High enzyme concentration may result in **Star Activity**.
- \* Total reaction volume dependent on experiment.
- \* The amount of enzyme to be used is very much dependent on the DNA template.
- \* For plasmid DNA, 5-10X more enzyme is required.

#### Example of Digestion Reaction

Enzyme : 1 unit  
Lambda 0.3μg/μl : 3.33μl (1μg DNA)  
10X Reaction Buffer : 5μl  
Sterile Distilled Water : Up to 50μl

#### Product Use Limitation

This product is for research purposes and *in vitro* use only.